

Student Name: _____

Score: _____

Derivatives using Power Rule

Find the derivatives using power rule:

$$y = 10x^3$$

$$y = \frac{1}{2}x^{-2}$$

$$y = \frac{1}{2\sqrt{x}}$$

$$y = 3x^{\frac{-1}{15}}$$

$$y = 8x^6 + 2x^{17}$$

$$y = \sqrt[5]{x}$$

$$y = x^{\frac{1}{31}} + x^{\frac{-1}{7}}$$

$$y = 2x^{12} + 6x^7 + x^4$$

$$y = \frac{5}{3}x^3 - \frac{7}{6}x^6 + \frac{6}{4}x^8$$

$$y = \frac{1}{2}x^{\frac{3}{2}} - \frac{22}{7}x^{\frac{-5}{2}} + x^{\frac{3}{7}}$$

Student Name: _____

Score: _____

Derivatives using Power Rule

Find the derivatives using power rule:

$$y = \frac{8x^5 + 4x^4}{2x^2}$$

$$y = \frac{15x^7 + 21x^5 + 12x^3}{3x}$$

$$y = \frac{-22x^{-5} - 17x^{-11}}{21x^{-4}}$$

$$y = \frac{2x^{\frac{11}{3}} + 4x^{\frac{5}{4}} - 3x^{\frac{7}{2}}}{4x^{\frac{2}{3}}}$$

$$y = \frac{7x^2 + 5x^9}{4x^7}$$

$$y = \frac{\sqrt{x} - \sqrt[3]{x}}{\sqrt[5]{x}}$$

$$y = \frac{5x^{-45} + 15x^{-4} - 5x^{-17}}{5x^{-2}}$$

$$y = \frac{5x^2 + 12x^{-5}}{\sqrt{x}}$$

$$y = \frac{\frac{2}{7}x^{\frac{-5}{11}} + \frac{16}{7}x^{\frac{-12}{11}}}{x^{\frac{-21}{11}}}$$

$$y = \frac{x^{\frac{7}{3}} + x^{\frac{10}{3}}}{\sqrt[3]{x}}$$

Student Name: _____

Score: _____

Derivatives using Product Rule

Find the derivatives using product rule:

$$y = (x^2 + 1)(x + 1)^2$$

$$y = (x + 1)(\sqrt{x} + 2)$$

$$y = (x^2 + x + 1)(x - 1)$$

$$y = x(x^7 + 15)^3$$

$$y = x^2(x + 7)^3$$

$$y = x^7 \sqrt{4x^2 + 7}$$

$$y = \sqrt{x}(x^2 + 4)$$

$$y = (\sqrt{x^2 + 1})(\sqrt{x^2 - 1})$$

$$y = (2x + 1)(2x - 1)^4$$

$$y = (\sqrt{x} - 1)(\sqrt{x} + 1)$$

Student Name: _____

Score: _____

Derivatives using Quotient Rule

Find the derivatives using quotient rule:

$$y = \frac{x}{x+1}$$

$$y = \frac{x^2}{3x-1}$$

$$y = \frac{4x^3 - 7x}{5x^2 + 2}$$

$$y = \frac{(x^2 - 1)^3}{x^2 + 1}$$

$$y = \frac{x^9 - 1}{\sqrt{x^2 - 1}}$$

$$y = \frac{4x^2}{x^3 + 3}$$

$$y = \frac{x^3}{\sqrt{x} + 1}$$

$$y = \frac{x+1}{x-1}$$

$$y = \frac{x^2 + 3}{x^2 + 4}$$

$$y = \frac{\sqrt{x}}{x + \frac{7}{2}}$$

Techniques of Differentiation - Classwork

Taking derivatives is a process that is vital in calculus. In order to take derivatives, there are rules that will make the process simpler than having to use the definition of the derivative.

1. The constant rule: The derivative of a constant function is 0. That is, if c is a real number, then $\frac{d}{dx}[c] = 0$.

a) $y = 7$

$y' =$

b) $f(x) = 0$

$f'(x) =$

c) $s(t) = -8$

$s'(t) =$

d) $y = \pi x^3$

$\frac{dy}{dx} =$

2. The single variable rule: The derivative of x is 1. $\frac{d}{dx}[x] = 1$. This is consistent with the fact that the slope of the line $y = x$ is 1.

a) $y = x$

$y' =$

b) $f(x) = x$

$f'(x) =$

c) $s(t) = t$

$s'(t) =$

3. The power rule: If n is a rational number then the function x^n is differentiable and $\frac{d}{dx}[x^n] = nx^{n-1}$.

Take the derivatives of the following. Use correct notation.

a) $y = x^2$

b) $f(x) = x^6$

c) $s(t) = t^{30}$

d) $y = \sqrt{x}$

e) $y = \frac{1}{x}$

f) $f(x) = \frac{1}{x^3}$

g) $s(t) = \frac{1}{\sqrt[3]{t}}$

h) $y = \frac{1}{x^{3/4}}$

4) The constant multiple rule: If f is a differentiable function and c is a real number, then $\frac{d}{dx}[cf(x)] = cf'(x)$

Take the derivatives of the following. Use correct notation.

a) $y = \frac{2}{x^2}$

b) $f(x) = \frac{4x^3}{3}$

c) $s(t) = -t^5$

d) $y = 4\sqrt{x}$

e) $y = \frac{-5}{3x^3}$

f) $f(x) = \frac{-5}{(3x)^3}$

g) $s(t) = \frac{4}{\sqrt{t}}$

h) $y = \frac{-12}{\sqrt[3]{x^5}}$

5. The sum and difference rules. The derivative of a sum or difference is the sum or difference of the derivatives.

$$\frac{d}{dx}[f(x) + g(x)] = f'(x) + g'(x) \quad \text{and} \quad \frac{d}{dx}[f(x) - g(x)] = f'(x) - g'(x)$$

Take the derivatives of the following. Use correct notation.

a) $y = x^2 + 5x - 3$

b) $f(x) = x^4 - \frac{3}{2}x^3 + 2x^2 + x - 6$

c) $y = \frac{4}{x} - \frac{4}{x^2} + \frac{4}{x^3}$

d) $y = 6\sqrt{x}(\sqrt[3]{x} - 2x + 6)$

e) $f(x) = (2x - 3)^2$

f) $y = (x^2 - x + 1)^2$

6. **The Product Rule:** The derivative of the product of two functions is the first times the derivative of the second

plus the second times the derivative of the first. $\frac{d}{dx}[f(x) \cdot g(x)] = f(x) \cdot g'(x) + g(x) \cdot f'(x)$

a) Find y' if $y = (4x - 2x^2)(3x - 5)$

without product rule

b) Find y' if $y = (x^2 - x + 1)^2$

with product rule

c) Find $f'(x)$ if $f(x) = (3x^2 - 2x + 5)(-5x^4 + 2x^3 - 7x^2 + x + 2)$

7. **The Quotient Rule:** The derivative of the quotient of two functions f and g can be found using the following:

$$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{g(x) \cdot f'(x) - f(x) \cdot g'(x)}{[g(x)]^2}$$

a) Find $\frac{d}{dx}\left[\frac{5x+2}{x^2-1}\right]$

b) Find $\frac{d}{dx}\left[\frac{5x+3}{x^2+4x-2}\right]$

c) Find $\frac{d}{dx}\left[\frac{x^3 - 4x^2 + 4x - 2}{2x}\right]$

d) Find $\frac{d}{dx}\left[\frac{2x}{x^3 - 4x^2 + 4x - 2}\right]$

Techniques of Differentiation - Homework

For the following functions, find $f'(x)$ and $f'(c)$ at the indicated value of c .

$$1) f(x) = x^2 - 6x + 1 \quad c = 0 \quad 2) f(x) = \frac{1}{x} - \frac{3}{x^2} + \frac{4}{x^3} \quad c = 1 \quad 3) f(x) = 3\sqrt{x} - \frac{1}{\sqrt[3]{x}} \quad c = 1$$

For the following functions, find the derivative using the power rule.

$$4) y = \frac{8}{3x^2}$$

$$5) y = \frac{-9}{(3x^2)^3}$$

$$6) y = \frac{6x^{3/2}}{x}$$

$$7) y = \frac{4x^2 - 5x + 6}{3}$$

$$8) y = \frac{x^2 - 6x + 2}{2x}$$

$$9) y = \frac{x^3 + 8}{x + 2}$$

$$10) y = x^4 - \frac{3}{2}x^3 + 5x^2 - 6x - 2$$

$$11) y = \frac{x^3 - 3x^2 + 10x - 5}{x^2}$$

$$12) y = (x^2 + 4x)(2x - 1)$$

$$13) y = (x - 2)^3$$

$$14) y = \sqrt[3]{x} - \sqrt[3]{x^2}$$

$$15) y = \frac{(x^2 - x + 2)^2}{x}$$

For the following functions, find the derivatives.

$$16) y = (x^2 - 4x - 6)(x^3 - 5x^2 - 3x) \quad 17) y = \frac{3x - 2}{2x + 3}$$

$$18) y = \frac{x^2 - 4x - 2}{x^2 - 1}$$

19) $y = \frac{x-1}{\sqrt{x}}$

20) $y = \frac{x^2 - x + 1}{\sqrt[3]{x}}$

21) $y = \left(\frac{x-3}{x+4}\right)(3x-2)$

22) $y = \frac{x-1}{x^2 + 2x + 2}$

23) $y = \frac{x^2 + k^2}{x^2 - k^2}$, k is a constant

24) $y = \frac{x^2 - k^2}{x^2 + k^2}$, k a constant

Find an equation of the tangent line to the graph of f at the indicated point and then use your calculator to confirm the results.

25) $f(x) = \frac{x^2}{x-1}$ at $(2, 4)$

26) $f(x) = (x-2)(x^2 - 3x - 1)$ at $(-1, -9)$

27) $f(x) = \frac{x^2 - 4x + 2}{2x - 1}$ at $\left(2, -\frac{2}{3}\right)$

28) $y = \left(\frac{x+3}{x+1}\right)(4x+1)$ at $\left(-\frac{1}{2}, -5\right)$